

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/41	A1	(11) International Publication Number: WO 96/40119 (43) International Publication Date: 19 December 1996 (19.12.96)
<p>(21) International Application Number: PCT/US96/07444</p> <p>(22) International Filing Date: 22 May 1996 (22.05.96)</p> <p>(30) Priority Data: 08/473,819 7 June 1995 (07.06.95) US</p> <p>(71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).</p> <p>(72) Inventor: CAMDEN, James, Berger; 7339 Charter Cup Lane, West Chester, OH 45069 (US).</p> <p>(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report.</i></p>
<p>(54) Title: USE OF 1,2,4-TRIAZOLE DERIVATIVES FOR THE MANUFACTURE OF A MEDICAMENT FOR THE TREATMENT OF CANCERS</p> <p>(57) Abstract</p> <p>A pharmaceutical composition that inhibits the growth of tumors and cancers in mammals that comprises a material is disclosed. The particular material used is a 1H-1,2,4-triazole derivative. These compounds can also be used to treat viral infections.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

USE OF 1,2,4-TRIAZOLE DERIVATIVES FOR THE MANUFACTURE OF A MEDICAMENT FOR THE TREATMENT OF CANCERS

TECHNICAL FIELD

5 This invention is a pharmaceutical composition that inhibits the growth of cancers, leukemia and tumors in mammals, particularly in human and warm blooded animals. The composition contains a 1H-1,2,4-triazole derivative. The compositions can also be used to treat viral infections.

BACKGROUND OF THE INVENTION

10 Cancers are the leading cause of death in animals and humans. The exact cause of cancer is not known, but links between certain activities such as smoking or exposure to carcinogens and the incidence of certain types of cancers and tumors has been shown by a number of researchers.

15 Many types of chemotherapeutic agents have been shown to be effective against cancers and tumor cells, but not all types of cancers and tumors respond to these agents. Unfortunately, many of these agents also destroy normal cells. The exact mechanism for the action of these chemotherapeutic agents are not always known.

20 Despite advances in the field of cancer treatment the leading therapies to date are surgery, radiation and chemotherapy. Chemotherapeutic approaches are said to fight cancers that are metastasized or ones that are particularly aggressive. Such cytotoxic or cytostatic agents work best on cancers with large growth factors, i.e., ones whose cells are rapidly dividing. To date, hormones, in particular estrogen, progesterone and testosterone, and some antibiotics produced by a variety of
25 microbes, alkylating agents, and anti-metabolites form the bulk of therapies available to oncologists. Ideally cytotoxic agents that have specificity for cancer and tumor cells while not affecting normal cells would be extremely desirable. Unfortunately, none have been found and instead agents which target especially rapidly dividing cells (both tumor and normal) have been used.

30 Clearly, the development of materials that would target tumor cells due to some unique specificity for them would be a breakthrough. Alternatively, materials that were cytotoxic to tumor cells while exerting mild effects on normal cells would be desirable. Therefore, it is an object of this invention to provide a pharmaceutical composition that is effective in inhibiting the growth of tumors and cancers in
35 mammals with mild or no effects on normal cells.


More specifically, it is an object of this invention to provide an anti-cancer composition comprising a pharmaceutical carrier and a 1H-1,2,4-triazole derivative as

defined herein along with a method for treating such cancers.

These and other objects will become evident from the following detailed description of this inventions.

SUMMARY OF THE INVENTION

- 5 A pharmaceutical composition for treatment of mammals, and in particular, warm blooded animals and humans, comprising a pharmaceutical carrier and an effective amount anti-cancer compound selected from the group consisting of:



- wherein Z is an alkylene selected from the group consisting of
 10 $\text{CH}_2\text{-CH}_2\text{-}$, $\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-}$, $\text{-CH(CH}_3\text{)-CH(CH}_3\text{)-}$ and $\text{-CH}_2\text{-CH(alkyl)-}$ wherein said alkyl has from 1 to about 10 carbon atoms; and Ar is a member selected from the group consisting of phenyl, substituted phenyl, thienyl, halothienyl, naphthyl and fluorenyl, wherein "substituted phenyl" has the meaning of a phenyl radical having
 15 thereon from 1 to 3 substituents selected independently from the group consisting of halo, lower alkyl, lower alkyloxy, cyano and nitro. The therapeutically active acid addition salts of the foregoing compound (I) are also embraced within the scope of this invention.

- As used in the foregoing definition of Z, the term "alkyl" is meant to include straight and branch chained hydrocarbon radicals having from 1 to about 10 carbon
 20 atoms, such as, for example, methyl, ethyl, 1-methylethyl, propyl, 1,1-dimethylethyl, butyl, pentyl, hexyl, heptyl, octyl, decyl and the like; as used herein "lower alkyl" may be straight or branch chained saturated hydrocarbons having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, propyl, 1-methylethyl, butyl, 1,1-dimethylethyl, pentyl, hexyl and the like alkyls; and the term "halo" is generic to halogen atoms of
 25 atomic weight less than 127; i.e., fluoro, chloro, bromo and iodo.

- These compositions can be used to inhibit the growth of cancers and other tumors in humans or animals by administration of an effective amount either orally, rectally, topically or parenterally, intravenously or by injection into the tumor. These compositions do not significantly affect healthy cells as compared to adriamycin which
 30 has a detrimental effect on healthy cells.

These compositions are also be used to treat viruses

DETAILED DESCRIPTION OF THE INVENTION

A. Definitions:

As used herein, the term "comprising" means various components can be conjointly employed in the pharmaceutical composition of this invention. Accordingly,
5 the terms "consisting essentially of" and "consisting of" are embodied in the term comprising.

As used herein, a "pharmaceutically acceptable" component is one that is suitable for use with humans and/or animals without undue adverse side effects (such as toxicity, irritation, and allergic response) commensurate with a reasonable
10 benefit/risk ratio.

As used herein, the term "safe and effective amount" refers to the quantity of a component which is sufficient to yield a desired therapeutic response without undue adverse side effects (such as toxicity, irritation, or allergic response) commensurate with a reasonable benefit/risk ratio when used in the manner of this invention. The
15 specific "safe and effective amount" will, obviously, vary with such factors as the particular condition being treated, the physical condition of the patient, the type of mammal being treated, the duration of the treatment, the nature of concurrent therapy (if any), and the specific formulations employed and the structure of the compounds or its derivatives.

As used herein, a "pharmaceutical addition salts" is salt of the anti-cancer compound with an organic or inorganic acid. These preferred acid addition salts are chlorides, bromides, sulfates, nitrates, phosphates, sulfonates, formates, tartrates, maleates, malates, citrates, benzoates, salicylates, ascorbates, and the like.
20

As used herein, a "pharmaceutical carrier" is a pharmaceutically acceptable
25 solvent, suspending agent or vehicle for delivering the anti-cancer agent to the animal or human. The carrier may be liquid or solid and is selected with the planned manner of administration in mind.


As used herein, "cancer" refers to all types of cancers or neoplasm or malignant tumors and all types of cancers including leukemia that are found in mammals.

As used herein, the "anti-cancer compounds" are the 1H-1,2,4-triazoles and their salts. The exact 1H-1,2,4-triazoles are described in detail below. The preferred materials are the products sold under the names "propiconazole®" by Janssen
30 Pharmaceutica NV (Belgium).

As used herein, "viruses" includes viruses which cause diseases (viral
35 infections) in man and other warm blooded animals, such as HIV virus, herpes, influenza and rhinoviruses.

B. THE ANTI-CANCER COMPOUNDS

The anti-cancer compounds are 1H-1,2,4-triazole derivatives which are known for their antifungal activities. They are systemic materials used to prevent and eradicate fungi. The compounds have the following structure:



5

wherein Z is an alkylene selected from the group consisting of CH₂-CH₂-, -CH₂-CH₂-CH₂-, -CH(CH₃)-CH(CH₃)- and -CH₂-CH(alkyl) wherein said alkyl has from 1 to about 10 carbon atoms; and Ar is a member selected from the group consisting of phenyl, substituted phenyl, thienyl, halothienyl, naphthyl and fluorenyl, wherein "substituted phenyl" has the meaning of a phenyl radical having thereon from 1 to 3 substituents selected independently from the group consisting of halo, lower alkyl, lower alkyloxy, cyano and nitro. The therapeutically active acid addition salts of the foregoing compound (I) are also embraced within the scope of this invention.

15

As used in the foregoing definition of Z, the term "alkyl" is meant to include straight and branch chained hydrocarbon radicals having from 1 to about 10 carbon atoms, such as, for example, methyl, ethyl, 1-methylethyl, propyl, 1,1-dimethylethyl, butyl, pentyl, hexyl, heptyl, octyl, decyl and the like; as used herein "lower alkyl" may be straight or branch chained saturated hydrocarbons having from 1 to 6 carbon atoms, such as, for example, methyl, ethyl, propyl, 1-methylethyl, butyl, 1,1-dimethylethyl, pentyl, hexyl and the like alkyls; and the term "halo" is generic to halogen atoms of atomic weight less than 127; i.e., fluoro, chloro, bromo and iodo.

20

Their pharmaceutically acceptable acid addition salts with both organic and inorganic acids can also be used herein.

25

Preferred derivatives include:

- 1-[2-(2,4-dichlorophenyl)-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole;
 - 1-[2-(2,4-dichlorophenyl)-4-methyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 - 1-[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 - 1-[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 - 1-[2-(2,4-dichlorophenyl)-4-pentyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole, and
- the therapeutically active acid addition salts thereof.

30

These compounds are prepared according to the method described in U.S. 4,079,062 issued to Van Reet. et al, Mar 14, 1978.

It is believed that these particular materials have the capability of reducing tumors or decreasing their growth significantly because of their ability to inhibit the synthesis of sterols.

C. DOSAGE

5 Any suitable dosage may be given in the method of the invention. The type of disease (cancer, leukemia or virus), the compound, the carrier and the amount will vary widely depending on the species of the warm blooded animal or human, body weight, and tumor being treated. Generally a dosage of between about 2 milligrams (mg) per kilogram (kg) of body weight and about 400 mg per kg of body weight is
10 suitable. Preferably from 15 mg to about 150 mg/kg of body weight is used. Generally, the dosage in man is lower than for small warm blooded mammals such as mice. A dosage unit may comprise a single compound or mixtures thereof with other compounds or other cancer inhibiting compounds. The dosage unit can also comprise diluents, extenders, carriers and the like. The unit may be in solid or gel form such as
15 pills, tablets, capsules and the like or in liquid form suitable for oral, rectal, topical, intravenous injection or parenteral administration or injection into or around the tumor.

D. DOSAGE DELIVERY FORMS

The anti-cancer compounds are typically mixed with a pharmaceutically
20 acceptable carrier. This carrier can be a solid or liquid and the type is generally chosen based on the type of administration being used. The active agent can be coadministered in the form of a tablet or capsule, as an agglomerated powder or in a liquid form. Examples of suitable solid carriers include lactose, sucrose, gelatin and agar. Capsule or tablets can be easily formulated and can be made easy to swallow or
25 chew; other solid forms include granules, and bulk powders. Tablets may contain suitable binders, lubricants, diluents, disintegrating agents, coloring agents, flavoring agents, flow-inducing agents, and melting agents. Examples of suitable liquid dosage forms include solutions or suspensions in water, pharmaceutically acceptable fats and oils, alcohols or other organic solvents, including esters, emulsions, syrups or elixirs,
30 suspensions, solutions and/or suspensions reconstituted from non-effervescent granules and effervescent preparations reconstituted from effervescent granules. Such liquid dosage forms may contain, for example, suitable solvents, preservatives, emulsifying agents, suspending agents, diluents, sweeteners, thickeners, and melting agents. Oral dosage forms optionally contain flavorants and coloring agents. Parenteral and
35 intravenous forms would also include minerals and other materials to make them compatible with the type of injection or delivery system chosen.

Specific examples of pharmaceutical acceptable carriers and excipients that

may be used to formulate oral dosage forms of the present invention are described in US. Pat. No. 3,903,297 to Robert, issued Sept. 2, 1975. Techniques and compositions for making dosage forms useful in the present invention are described in the following references: 7 Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, Editors, 1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms 2nd Edition (1976).

E. METHOD OF TREATMENT

The method of treatment can be any suitable method which is effective in the treatment of the particular cancer type or virus that is being treated. Treatment may be oral, rectal, topical, parenteral or intravenous administration or by injection into the tumor and the like. The method of applying an effective amount also varies depending on the tumor being treated. It is believed that parenteral treatment by intravenous, subcutaneous, or intramuscular application of the 1H-1,2,4-triazole compounds, formulated with an appropriate carrier, additional cancer inhibiting compound or compounds or diluent to facilitate application will be the preferred method of administering the compounds to warm blooded animals.

The method of treating viral infections may also be by oral, rectal, topical, parenteral or intravenous administration.

In vitro Data

The following examples are illustrative and are not meant to be limiting to the invention.

Colon, Breast and Lung Tumor Cells Test

The following cell culture tests were performed to test the toxicity of the N-phosphonoglycine compounds on colon, breast and lung human tumor cells. The viability of the cells were tested by looking at MTT (3-[4,5-dimethylthiazol-2-yl] -2,5-diphenyltetrazolium bromide) reduction. MTT assay is a well known measure of cell viability.

The colon tumor cells (HT29 from American Type Culture Collection (ATCC)) and the breast cells (MX1 from cell lines from ATCC) were cultured in Eagle's Minimal Essential Medium with 10% fetal bovine serum. The lung tumor cells (A549 from ATCC cell lines) were cultured in Ham's F12 medium with 10% fetal bovine serum.

The tumor cells were passaged and seeded into culture flasks at the desired cell densities. The culture medium was decanted and the cell sheets were washed twice with phosphate buffered saline (PBS). The cells were trypsinized and triturated prior to seeding the flasks. Unless otherwise indicated the cultures were incubated at $37 \pm 1^\circ \text{C}$ in a humidified atmosphere of $5 \pm 1\%$ carbon dioxide in air. The cultures were

incubated until they were 50-80% confluent.

The cells were subcultured when the flasks were subconfluent. The medium was aspirated from the flasks and the cell sheets rinsed twice with PBS. Next, the Trypsin Solution was added to each flask to cover the cell sheet. The Trypsin Solution was removed after 30-60 seconds and the flasks were incubated at room temperature for two to six minutes. When 90% of the cells became dislodged, growth medium was added. The cells were removed by trituration and transferred to a sterile centrifuge tube. The concentration of cells in the suspension was determined, and an appropriate dilution was made to obtain a density of 5000 cells/ml. The cells were subcultured into the designated wells of the 96-well bioassay plates (200 microliter cell suspension per well). PBS was added to all the remaining wells to maintain humidity. The plates were then incubated overnight before test article treatment.

Each dose of test article was tested by treating quadruplicate wells of cultures with 100 microliter of each dilution. Those wells designated as solvent controls received an additional 100 microliter of methanol control; negative controls wells received an additional 100 microliters of treatment medium. PBS was added to the remaining wells not treated with test article or medium. The plates were then incubated for approximately 5 days.

At the end of the 5 day incubation, each dose group was examined microscopically to assess toxicity. A 0.5 mg/ml dilution of MTT was made in treatment medium, and the dilution was filtered through a 0.45 micrometer filter to remove undissolved crystals. The medium was decanted from the wells of the bioassay plates. Immediately thereafter, 2000 microliter of the filtered MTT solution was added to all test wells except for the two untreated blank test wells. The two blank wells received 200 microliters of treatment medium. The plates were returned to the incubator for about 3 hours. After incubation, the MTT containing medium was decanted. Excess medium was added to each well and the plates were shaken at room temperature for about 2 hours.

The absorbance at 550 nm (OD₅₅₀) of each well was measured with a Molecular Devices (Menlo Park, CA) VMax plate reader.

The mean OD₅₅₀ of the solvent control wells and that of each test article dilution, and that of each of the blank wells and the positive control were calculated. The mean OD₅₅₀ of the blank wells was subtracted from the mean of the solvent control wells, and test article wells, respectively to give the corresponding mean OD₅₅₀.

$$\% \text{ of Control} = \frac{\text{corrected mean OD}_{550} \text{ of Test Article Dilution}}{\text{corrected mean of OD}_{550} \text{ of Solvent Control}} \times 100$$

Dose response curves were prepared as semi-log plots with % of control on the ordinate (linear) and the test article concentration on the abscissa (logarithmic). The EC₅₀ was interpolated from the plots for each test article.

For the test articles administered in methanol, separate responses were prepared to correct for the methanol data.


Adriamycin was used as a positive control. In all cases, it was more toxic than any of the test materials by one or two logs. Adriamycin is one of the more potent agents in current use and one with significant side effects. The peak plasma concentration of other, quite effective chemotherapeutic agents may be 10 to 50 times higher than that of Adriamycin. The EC-50 is the concentration at which one half the cells are killed.

Table 1

<u>Test Material</u>	<u>EC-50 Result (ppml)</u>		
	HT29	MX1	A549
Adriamycin	0.00639	0.00078	0.00373
Propiconazole	0.0331	0.0284	0.113

These experiments show that these compositions are effective in killing tumor cells without significantly affecting healthy cells. They are safer than adriamycin.

1. A pharmaceutical composition for treating cancers and viruses comprising a safe and effective amount of:



wherein Z is an alkylene selected from the group consisting of $\text{CH}_2\text{-CH}_2\text{-}$, $\text{-CH}_2\text{-CH}_2\text{-CH}_2\text{-}$, $\text{-CH(CH}_3\text{)-CH(CH}_3\text{)-}$ and $\text{-CH}_2\text{-CH(alkyl)-}$ wherein said alkyl has from 1 to 10 carbon atoms; and Ar is a member selected from the group consisting of phenyl, substituted phenyl, thienyl, halothienyl, naphthyl and fluorenyl.

2. A pharmaceutical composition according to Claim 1 comprising a pharmaceutically acceptable carrier and a safe and effective amount of a 1H-1,2,4-triazole selected from the group consisting of:
 1-[2-(2,4-dichlorophenyl)-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole;
 1-[2-(2,4-dichlorophenyl)-4-methyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 1-[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 1-[2-(2,4-dichlorophenyl)-4-propyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole,
 1-[2-(2,4-dichlorophenyl)-4-pentyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole, and
 the therapeutically active acid addition salts thereof.
3. A pharmaceutical composition according to Claim 1 or 2 wherein said pharmaceutical acceptable acid addition salts are selected from the group consisting of chlorides, bromides, sulfates, nitrates, phosphates, sulfonates, formates, tartrates, maleates, malates, citrates, benzoates, salicylates, ascorbates and mixtures thereof.
4. A method of treating cancer in warm blooded mammals comprising administering 2 mg/kg body weight to 400 mg/kg of a 1H-1,2,4-triazole derivative according to Claims 1 or 2.
5. A method according to Claim 4 wherein said 1H-1,2,4-triazole is administered orally or enterically, intravenously, peritoneally, parenterally or by injection into the tumor.

6. A method according to Claim 4 wherein said 1H-1,2,4-triazole is administered in a solid form, wherein said solid form includes a carrier selected from the group consisting of lactose, sucrose, gelatin and agar.
7. A method according to Claim 4 wherein said 1H-1,2,4-triazole is administered in a liquid form, wherein said liquid dosage form is selected from the group consisting of aqueous solutions, alcohol solutions, emulsions, suspensions, and suspensions reconstituted from non-effervescent and effervescent preparations and suspensions in pharmaceutically acceptable fats or oils.
8. A unit dosage composition for treating cancer and viral infections in animals or humans comprising a 1H-1,2,4-triazole according to Claims 1 or 2.
9. A method of treating viral infections in warm blooded mammals comprising administering from 2 mg/kg to 400 mg/kg body weight of a 1H-1,2,4-triazole derivative according to Claims 1 or 2.

INTERNATIONAL SEARCH REPORT

International Application No.
PCT/US 96/07444

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/41

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 160 838 A (GUSTAAF VAN REET ET AL.) 10 July 1979 see column 4, line 5 - line 10; claims 1-7; example 1 ---	1-3,9
X	US 4 079 062 A (GUSTAAF VAN REET ET AL.) 14 March 1978 cited in the application see column 3, line 64 - column 4, line 3; claims 1-6 -----	1-3,9

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

*** Special categories of cited documents :**

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *A* document member of the same patent family

Date of the actual completion of the international search

24 July 1996

Date of mailing of the international search report

02.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patendaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl.
Fax (+ 31-70) 340-3016

Authorized officer

Tzschoppe, D

INTERNATIONAL SEARCH REPORT

ernational application No.

PCT/US 96/ 07444

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. Claims 1-9
2. Claim 10

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Claims 1-9

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/US 96/07444

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4160838	10-07-79	NONE	
US-A-4079062	14-03-78	AT-B- 347942	15-06-78
		AT-B- 351861	27-08-79
		AU-B- 503503	06-09-79
		BE-A- 835579	14-05-76
		CA-A- 1094079	20-01-81
		CH-A- 625103	15-09-81
		CH-A- 615676	15-02-80
		DE-A- 2551560	20-05-76
		FR-A- 2290898	11-06-76
		GB-A- 1522657	23-08-78
		NL-A- 7513389	20-05-76
		SE-B- 433495	28-05-84
		SE-B- 425246	13-09-82
		SE-A- 7512643	19-05-76
		SE-B- 433496	28-05-84
		SE-A- 8305128	22-09-83
		SI-A- 7512929	31-12-94

THIS PAGE BLANK (USPTO)